


Conserving Atlantic Bluefin Tuna with Spawning Sanctuaries



THE
PEW
ENVIRONMENT GROUP

SCIENCE BRIEF

Atantic bluefin tuna populations in both the western and eastern Atlantic Ocean are currently at very low levels (ICCAT 2010a; 2010b), and additional management measures are needed to rebuild their populations (Armsworth *et al.* 2010; Block *et al.* 2005; Druon 2010; Hurry *et al.* 2008; ICCAT 2007; MacKenzie *et al.* 2009; Safina and Klinger 2008; Teo *et al.* 2007a; Teo and Block 2010). The prohibition of certain fishing activities at specific times or in specific areas, known as “time and area closures” in fisheries management parlance, are tools commonly used to protect crucial genetic and biological diversity, restore population structure (e.g., age and sex distribution) and spawning stocks, and reduce bycatch (Pelletier *et al.* 2008).

Globally, pelagic fishes (Goodyear 1999), mollusks (Dredge 1992) and reef fishes (Galal *et al.* 2002) have been managed with closures because they protect specific size classes, sexes or individual species from excessive fishing mortality. For example, North Atlantic swordfish have been successfully protected by time and area closures and have recently been declared rebuilt by the U.S. National Marine Fisheries Service (NMFS 2009a), see box below. Recent research suggests that closure of spawning areas for Atlantic bluefin tuna, used specifically to protect spawning fish (Beets and Friedlander 1998; Heyman *et al.* 2005; Nemeth 2005; Sala *et al.* 2001), may be a viable tool to help rebuild their depleted populations (Armsworth *et al.* 2010; Block *et al.* 2005; Druon 2010; ICCAT 2007; Teo *et al.* 2007a; Teo and Block 2010).

The importance of genetic diversity

Genetic diversity is important for species survival because those species that have little genetic diversity may be more susceptible to disease or the effects of changes in the environment. Increased genetic diversity improves the chance of offspring with a variety of characteristics that could allow some of them to withstand changes in the environment and decreases the chance that deleterious genes (such as those linked to disease) show up in the population.

SPAWNING GROUND SANCTUARIES FOR ATLANTIC BLUEFIN TUNA

For spawning area closures to be implemented successfully, fisheries managers must have information on specific spawning times and locations, migration routes and habitat use behavior surrounding the spawning event (Nemeth *et al.* 2007). Atlantic bluefin tuna’s two known spawning locations are in the Mediterranean Sea (Fromentin and Powers 2005; ICCAT 2008b; Karakulak *et al.* 2004; Mather *et al.* 1995; Neilson and Campana 2008; Nishida *et al.* 1998; Rooker *et al.* 2007) and in the Gulf of Mexico (Mather *et al.* 1995). Within the Mediterranean Sea, scientists have identified six major regions of spawning activity, some of which may constitute genetically isolated populations (ICCAT 2010c).

Area closures rebuild swordfish populations

North Atlantic swordfish became overfished in the early 1990s, and by 1999 it had been determined that despite the implementation of a minimum size limit, juvenile swordfish were still suffering from a high fishing mortality rate, and this was likely prohibiting the population from recovering. Subsequently, the U.S. National Marine Fisheries Service instituted two permanent area closures to protect juvenile swordfish in U.S. waters. These closures prohibited pelagic longline fishing in areas in the Gulf of Mexico and off the east coast of Florida. By 2002, the population was considered to be in recovery and by 2009 was considered rebuilt, suggesting the use of area closures aided in the timely recovery of North Atlantic swordfish.

Conserving Atlantic Bluefin Tuna with Spawning Sanctuaries

In 2006, the Standing Committee on Research and Statistics (SCRS) of the International Commission for the Conservation of Atlantic Tunas (ICCAT), the intergovernmental organization charged with the management of tunas and tunalike species in the Atlantic Ocean and adjacent seas, recommended that the Mediterranean Sea be closed to fishing for Atlantic bluefin tuna during their spawning season. However, this scientific recommendation was not adopted by ICCAT itself (ICCAT 2006).

In 2008, ICCAT agreed to establish a multiyear recovery plan for Atlantic bluefin tuna in the eastern Atlantic Ocean and Mediterranean Sea, and the Commission requested that the SCRS identify possible spawning grounds in Mediterranean waters for the creation of sanctuaries that would aid in bluefin recovery (ICCAT 2008a). The Commission also upheld a recommendation to prohibit a directed bluefin tuna fishery in the Gulf of Mexico, where spawning areas are comparatively well known but not protected (ICCAT 2008a).

In addition, an independent review of ICCAT in 2008 recommended an immediate closure of all Atlantic bluefin tuna spawning grounds during their spawning seasons (Hurry *et al.* 2008). Despite these calls for action, spawning area sanctuaries have not been put into place. Recent genetic, tracking and other studies (Block *et al.* 2005; Boustany *et al.* 2008; Carlsson *et al.*

2004, 2007; Nemerson *et al.* 2000; Rooker *et al.* 2008a, 2008b) have added support to the idea that Atlantic bluefin tuna return to their birth locations to spawn and thus have genetically distinct populations. Research by Block *et al.* (2005) revealed two populations of Atlantic bluefin tuna that overlap on feeding grounds in the North Atlantic and subsequently migrate to separate spawning locations in the Gulf of Mexico and Mediterranean. Similarly, genetic analysis shows that populations from the Gulf of Mexico and eastern and western Mediterranean Sea (Carlsson *et al.* 2004) are genetically distinct (Boustany *et al.* 2008; Carlsson *et al.* 2007). These results indicate that establishing sanctuaries in both the Gulf of Mexico and the Mediterranean is crucial to ensuring that the genetic diversity of the species is preserved.

Gulf of Mexico

The U.S. National Marine Fisheries Service recently designated an area of the Gulf of Mexico as a Habitat of Particular Concern (HAPC) for spawning Atlantic bluefin tuna (NMFS 2009b), see Figure 1. However, the agency has not implemented time and area closures in this region that would protect spawning Atlantic bluefin tuna (NMFS 2006). While directed fishing of Atlantic bluefin tuna is not allowed in the Gulf (NMFS 2006), they are still incidentally caught (and large proportions subsequently die) by longlines targeting yellowfin tuna and swordfish (Armstrong *et al.* 2010; Teo and Block 2010).




Fig. 1 Distribution of known spawning areas for Atlantic bluefin tuna in the Gulf of Mexico

Teo and Block (2010) suggested that spatial and temporal management in the Gulf will become important in rebuilding the western Atlantic bluefin tuna populations. Their research showed that catch rates of Atlantic bluefin tuna in the Gulf increase dramatically during the breeding season (March to June) with peaks occurring in April and May. Block *et al.* (2005) suggested that considerable bycatch of Atlantic bluefin tuna in the Gulf of Mexico occurs during the spawning season and that this mortality could be substantially reduced by instituting time and area closures for longliners in the Gulf (Block *et al.* 2005).

Teo and Block (2010) provided additional information showing that targeted yellowfin tuna catch rates in the Gulf remain fairly constant throughout the year. Therefore, bycatch of bluefin tuna in this region can be substantially reduced by implementing time and area closures with only minimal impact on catch rates for target species (Teo and Block 2010). In addition, economic modeling has shown that time and area closures in the Gulf, combined with a coordinated effort among countries to rebuild bluefin tuna, would be economically beneficial over the long term, with an increase in profits of up to \$9 million a year (Armsworth *et al.* 2010).




Fig. 2 Distribution of known spawning areas for Atlantic bluefin tuna in the Mediterranean Sea

Mediterranean Sea

Atlantic bluefin tuna are found in the Mediterranean Sea throughout the year (Nemerson *et al.* 2000). Spawning begins in the eastern Mediterranean in May and moves west, ending in the Balearic Islands off the coast of Spain in July (Druon 2010; Heinisch *et al.* 2008; Karakulak *et al.* 2004). Spawning occurs offshore in the Balearic Sea and peak spawning densities occur around the Balearic Islands from the Mallorca Channel and south of Menorca (Alemany *et al.* 2010; Garcia *et al.* 2003, 2005), see Figure 2. A genetic and demographic analysis suggests that multiple distinct populations of Atlantic bluefin tuna may occur in the Mediterranean (Riccioni *et al.* 2010).

In the eastern Atlantic and Mediterranean Sea there are closures for large longliners and purse seine vessels, but there are no closures in effect that specifically address Atlantic bluefin tuna spawning grounds (Fromentin 2008). Druon (2010) suggested closing spawning areas in the Mediterranean during the spawning season while allowing fishing in restricted feeding grounds (areas where Atlantic bluefin tuna gather to feed) before and after the spawning closure. Additionally, Druon (2010) provided evidence that utilizing habitat mapping to determine open fishing areas might reduce the number of days at sea—and therefore fishing costs—used by fishermen to target and capture Atlantic bluefin tuna in this region.

SUMMARY

Time and area closures to protect spawning fish aggregations are a common management tool for protecting and restoring overexploited populations (Beets and Friedlander 1998; Heyman *et al.* 2005; Sala *et al.* 2001), and they have proved to be successful in rebuilding such overfished populations when their designations are based on scientifically accurate information (Nemeth 2005; Nemeth *et al.* 2007). Atlantic bluefin tuna spawning locations, spawning and habitat utilization behaviors, migration patterns and movement during spawning events are known (Block *et al.* 2005; Fromentin and Powers 2005; ICCAT 2008b; Nemerson *et al.* 2000; Nishida *et al.* 1998; Rooker *et al.* 2007; Teo *et al.* 2007a, 2007b), making them a viable candidate for spawning area closures.

Recent population assessments (ICCAT 2008b, 2010a, 2010b) indicate that Atlantic bluefin tuna populations in both the western and eastern Atlantic need increased protection to rebuild their depleted populations, yet current management measures are not working. According to this review and a number of experts (Armsworth *et al.* 2010; Block *et al.* 2005; Druon 2010; Hurry *et al.* 2008; ICCAT 2007; Teo *et al.* 2007a; Teo and Block 2010), spawning area closures could be an important option for restoring bluefin tuna populations.

References

- Alemaný, F., L. Quintanilla, P. Velez-Belchi, A. García, D. Cortés, J.M. Rodríguez, M.L. Fernández de Puelles, C. González-Pola and J.L. López-Jurado. 2010. Characterization of the spawning habitat of Atlantic bluefin tuna and related species in the Balearic Sea (western Mediterranean). *Progress in Oceanography* 86:21-38. Link
- Armsworth, P.R., B.A. Block, J. Eagle and J.E. Roughgarden. 2010. The economic efficiency of a time-area closure to protect spawning bluefin tuna. *Journal of Applied Ecology* 47:36-46. Link
- Beets, J., and A. Friedlander. 1998. Evaluation of a conservation strategy: A spawning aggregation closure for red hind, *Epinephelus guttatus*, in the US Virgin Islands. *Environmental Biology of Fishes* 55:91-98. Link
- Block, B.A., S.L.H. Teo, A. Walli, A. Boustany, M.J.W. Stokesbury, C.J. Farwell, K.C. Weng, H. Dewar and T.D. Williams. 2005. Electronic tagging and population structure of Atlantic bluefin tuna. *Nature* 434:1121-1127. Link
- Boustany, A.M., C.A. Reeb and B.A. Block. 2008. Mitochondrial DNA and electronic tracking reveal population structure of Atlantic bluefin tuna (*Thunnus thynnus*). *Marine Biology* 156:13-24. Link
- Carlsson, J., J.R. McDowell, P. Diaz-Jaimes, J.E.L. Carlsson, S.B. Boles, J.R. Gold and J.E. Graves. 2004. Microsatellite and mitochondrial DNA analyses of Atlantic bluefin tuna (*Thunnus thynnus thynnus*) population structure in the Mediterranean Sea. *Molecular Ecology* 13:3345-3356. Link
- Carlsson J., J.R. McDowell, J.E.L. Carlsson and J.E. Graves. 2007. Genetic identity of YOY bluefin tuna from the eastern and western Atlantic spawning areas. *Journal of Heredity* 98:23-28. Link
- Dredge, M.C.L. 1992. Using size limits to maintain scallop stocks in Queensland. In *Legal sizes and their use in fisheries management*, pp. 79-85. Australian Society for Fish Biology workshop. Lorne, Victoria, Australia. Link
- Druon, J. 2010. Habitat mapping of the Atlantic bluefin tuna derived from satellite data: Its potential as a tool for the sustainable management of pelagic fisheries. *Marine Policy* 34:293-297. Link
- Fromentin, J.M., and J.E. Powers. 2005. Atlantic bluefin tuna: population dynamics, ecology, fisheries and management. *Fish and Fisheries* 6:281-306. Link
- Fromentin, J.M. 2008. An attempt to evaluate the recent management regulations of the East Atlantic and Mediterranean bluefin tuna stock through a simple simulation model. *Collective Volume of Scientific Papers* 62:1271-1279. Link
- Galal, N., R.F.G. Ormond and O. Hassan. 2002. Effect of a network of no-take reserves in increasing catch per unit effort and stocks of exploited reef fish at Nabq, South Sinai, Egypt. *Marine and Freshwater Research* 53:199-205. Link
- García, A., F. Alemany, P. Velez-Belchi, J.M. Rodríguez, J.L. López Jurado, C. González Pola and J.M. de la Serna. 2003. Bluefin and frigate tuna spawning off the Balearic archipelago and the environmental conditions observed during the 2002 spawning season. *Collective Volume of Scientific Papers ICCAT* 55:1261-1270. Link
- García, A., F. Alemany, J.M. de la Serna, I. Oray, S. Karakulak, L. Rollandi, A. Arigò and S. Mazzola. 2005. Preliminary results of the 2004 bluefin tuna larval surveys off different Mediterranean sites (Balearic Archipelago, Levantine Sea and the Sicilian Channel). *Collective Volume of Scientific Papers ICCAT* 58:1261-1270. Link
- Goodyear, C.P. 1999. An analysis of the possible utility of time-area closures to minimize billfish bycatch by U.S. pelagic longlines. *Fishery Bulletin* 97:243-255. Link
- Heinisch, G., A. Corriero, A. Medina, F.J. Abascal, J. de la Serna, R. Vassallo-Agius, A. B. Ríos, A. García, F. de la Gándara, C. Fauvel, C.R. Bridges, C.C. Mylonas, S.F. Karakulak, I. Oray, G. De Metrio, H. Rosenfeld and H. Gordin. 2008. Spatial-temporal pattern of bluefin tuna gonad maturation across the Mediterranean Sea. *Marine Biology* 154:623-630. Link
- Heyman, W.D., B. Kjerfve, R.T. Graham, K.L. Rhodes and L. Garbutt. 2005. Spawning aggregations of *Lutjanus cyanopterus* (Cuvier) on the Belize Barrier Reef over a 6 year period. *Journal of Fish Biology* 67:83-101. Link
- Hurry, G.D., M. Hayashi and J.J. Maguire. 2008. Report of the independent review. International Commission for the Conservation of Atlantic Tunas. *International Commission for the Conservation of Atlantic Tunas* PLE-106/2000. Link
- International Commission for the Conservation of Atlantic Tunas. 2006. *Report for the biennial period, 2006-07*. Part 1 (2006)-Vol. 2 SCRS. Link
- International Commission for the Conservation of Atlantic Tunas. 2007. Species executive summary: Bluefin tuna. In *Report for the biennial period, 2006-07*. Part II (2007)-Vol. 2-SCRS, pp 98-114.
- International Commission for the Conservation of Atlantic Tunas. 2008a. *Report for biennial period, 2008-09*. Part 1 (2008) – Vol. 1, COM. Link
- International Commission for the Conservation of Atlantic Tunas. 2008b. *Report of the 2008 Atlantic bluefin tuna stock assessment session*. Link
- International Commission for the Conservation of Atlantic Tunas. 2010a. *Scientific Committee on Research and Statistics, bluefin tuna western Atlantic executive summary*. ICCAT Document Number SCI-019/2010.
- International Commission for the Conservation of Atlantic Tunas. 2010b. *Scientific Committee on Research and Statistics, bluefin tuna eastern Atlantic and Mediterranean Sea executive summary*. ICCAT Document Number SCI-018A/2010.
- International Commission for the Conservation of Atlantic Tunas. 2010c. *Response on bluefin tuna spawning areas*. ICCAT Document Number SCI-053A/2010.
- Karakulak, S., I. Oray, A. Corriero, A. Aprea, D. Spedicato, D. Zubani, N. Santamaria and G. De Metrio. 2004. First information on the reproductive biology of the bluefin tuna (*Thunnus thynnus*) in the eastern Mediterranean. *Collective Volume of Scientific Papers ICCAT* 56:1158-1162. Link
- MacKenzie, B.R., H. Mosegaard and A. Rosenberg. 2009. Impending collapse of bluefin tuna in the Northeast Atlantic and Mediterranean. *Conservation Letters* 2:26-35. Link
- Mather, F.J. III, J.M. Mason Jr. and A.C. Jones. 1995. Historical document: Life history and fisheries of Atlantic bluefin tuna. NOAA Technical Memorandum NMFS-SEFSC-370. Link
- National Marine Fisheries Service. 2006. *Final consolidated Atlantic highly migratory species fishery management plan*. NMFS Office of Sustainable Fisheries, Silver Spring, Md. Link
- National Marine Fisheries Service. 2009a. *Final Amendment 1 to the consolidated Atlantic highly migratory species fishery management plan essential fish habitat, including: a final environmental impact statement*. Highly Migratory Species Management Division, NMFS Office of Sustainable Fisheries, Silver Spring, Md. Link
- National Marine Fisheries Service. 2009b. *Stock Assessment and Fishery Evaluation (SAFE) report for Atlantic highly migratory species*. Highly Migratory Species Management Division, NMFS Office of Sustainable Fisheries, Silver Spring, Md. Link
- Neilson, J.D., and S.E. Campana. 2008. A validated description of age and growth of western Atlantic bluefin tuna (*Thunnus thynnus*). *Canadian Journal of Fisheries and Aquatic Sciences* 65:1523-1527. Link
- Nemerson D., S. Berkeley and C. Safina. 2000. Spawning site fidelity in Atlantic bluefin tuna, *Thunnus thynnus*: The use of size-frequency analysis to test for the presence of migrant East Atlantic bluefin tuna on Gulf of Mexico spawning grounds. *Fishery Bulletin* 98: 118-126. Link
- Nemeth, R.S. 2005. Population characteristics of a recovering US Virgin Islands red hind spawning aggregation following protection. *Marine Ecology Progress Series* 286:81-97. Link
- Nemeth, R.S., J. Blondeau, S. Herzlieb and E. Kadison. 2007. Spatial and temporal patterns of movement and migration at spawning aggregations of red hind, *Epinephelus guttatus*, in the US Virgin Islands. *Environmental Biology of Fishes* 78:365-381. Link
- Nishida, T., S. Tsuji and K. Segawa. 1998. Spatial data analyses of Atlantic bluefin tuna larval surveys in the 1994 ICCAT BYP. *Collective Volume of Scientific Papers ICCAT* 48:107-110. Link
- Pelletier, D., J. Claudet, J. Ferraris, L. Benedetti-Cecchi and J.A. Garcia-Charton. 2008. Models and indicators for assessing conservation and fisheries-related effects of marine protected areas. *Canadian Journal of Fisheries and Aquatic Sciences* 65: 765-779. Link
- Riccioni, G., M. Landi, G. Ferrara, I. Milano, A. Cariani, L. Zane, M. Sella, G. Barbujani and F. Tinti. 2010. Spatio-temporal population structuring and genetic diversity retention in depleted Atlantic bluefin tuna of the Mediterranean Sea. *PNAS* 107:2102-2107. Link
- Rooker, J.R., J.R. Alvarado Bremer, B.A. Block, H. Dewar, G. de Metrio, A. Corriero, R.T. Krause, E.D. Prince, E. Rodriguez-Marin and D.H. Secor. 2007. Life history and stock structure of Atlantic bluefin tuna (*Thunnus thynnus*). *Reviews in Fisheries Science* 15:265-310. Link
- Rooker, J.R., D.H. Secor, G. DeMetrio, A.J. Kaufman, A.B. Rios and V. Ticina. 2008a. Evidence of trans-Atlantic movement and natal homing of bluefin tuna from stable isotopes in otoliths. *Marine Ecology Progress Series* 368:231-239. Link
- Rooker, J.R., D.H. Secor, G. De Metrio, R. Schloesser, B.A. Block and J.D. Neilson. 2008b. Natal homing and connectivity in Atlantic bluefin tuna populations. *Science* 322: 742-744. Link
- Safina, C., and D. Klinger. 2008. Collapse of bluefin tuna in the western Atlantic. *Conservation Biology* 22:243-246. Link
- Sala, E., W. Ballesteros and R.M. Starr. 2001. Rapid decline of Nassau grouper spawning aggregations in Belize: Fishery management and conservation needs. *Fisheries* 26:23-30. Link
- Teo, S., A.M. Boustany and B.A. Block. 2007a. Oceanographic preferences of Atlantic bluefin tuna, *Thunnus thynnus*, on their Gulf of Mexico breeding grounds. *Marine Biology* 152:1105-1119. Link
- Teo, S., A.M. Boustany, H. Dewar, M.J.W. Stokesbury, K.C. Weng, S. Beemer, A.C. Setiz, C.J. Farwell, E.D. Prince and B.A. Block. 2007b. Annual migrations, diving behavior, and thermal biology of Atlantic bluefin tuna, *Thunnus thynnus*, on their Gulf of Mexico breeding grounds. *Marine Biology* 151:1-18. Link
- Teo, S., and B.A. Block. 2010. Comparative influence of ocean conditions on yellowfin and Atlantic bluefin tuna catch from longlines in the Gulf of Mexico. *PlosOne* 5:1-11. Link

The Pew Environment Group is the conservation arm of The Pew Charitable Trusts, a nongovernmental organization based in the United States, which applies a rigorous, analytical approach to improve public policy, inform the public and stimulate civic life.

901 E St. NW, 10th Floor,
Washington, D.C. 20004
Phone: 202-552-2000



E: oceanscience@PewTrusts.org www.PewEnvironment.org